

### **3.033 Variation in atmospheric CO<sub>2</sub> and CO concentrations emitted from biomass burning inferred from GOSAT/TANSO–FTS and MOPITT data.**

Early Career Scientist

Presenting Author:

**SHINGO KOSAKA**, Center for Environmental Remote Sensing (CEReS), Chiba University, Japan, [s.kosaka@chiba-u.jp](mailto:s.kosaka@chiba-u.jp)

Co-Authors:

**Naoko Saitoh**, Center for Environmental Remote Sensing (CEReS), Chiba University, Japan

Abstract:

This study has focused on biomass burning in Africa and studied the relationship between carbon monoxide (CO) and carbon dioxide (CO<sub>2</sub>) concentrations emitted due to the biomass burning by using CO<sub>2</sub> vertical profiles (Version 1) from the thermal infrared (TIR) band of Thermal and Near infrared Sensor for Carbon Observation Fourier Transform Spectrometer (TANSO–FTS) on board Greenhouse Gases Observing Satellite (GOSAT) [Saitoh et al., 2016] and CO vertical profiles (Version 6) retrieved from the combined use of the TIR and near infrared (NIR) bands of Measurement Of Pollution In The Troposphere (MOPITT) [Deeter et al., 2014].

In the latitude region around 10°N over Africa, TANSO–FTS CO<sub>2</sub> concentrations at 200–300 hPa were 2 ppm higher than a priori CO<sub>2</sub> concentrations taken from NIES–TM05 model [Saeki et al., 2013] from December to February when active biomass burning events occurred there frequently. MOPITT CO concentrations there were clearly higher in the lower troposphere from December to January and in the upper troposphere from February to March. We have compared monthly-averaged CO data from the MOPITT TIR/NIR bands with monthly-averaged detrended CO<sub>2</sub> data from the TANSO–FTS TIR band at 200–300 hPa in the northern low latitudes over Africa from 2009 to 2014. The comparison result showed that both TANSO–FTS CO<sub>2</sub> and MOPITT CO concentrations became higher almost at the same time. The periods of increase in the upper tropospheric CO<sub>2</sub> and CO concentrations were in January–March and in February–April, respectively, which slightly shifted from the periods of increase in their a priori concentrations. The amounts of the observed increase were larger compared to their a priori concentrations. We have analyzed backward trajectories of air masses with high CO and CO<sub>2</sub> concentrations observed by GOSAT/TANSO–FTS TIR band and MOPITT TIR/NIR bands to identify their sources.