

4.039 Tropospheric column ozone variability from space: results from the first multi-instrument intercomparison .

Presenting Author:

Audrey Gaudel, Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado; NOAA Earth System Research Laboratory, Chemical Sciences Division, audrey.gaudel@noaa.gov

Co-Authors:

Owen Cooper, Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado; NOAA Earth System Research Laboratory, Chemical Sciences Division

Valerie Thouret, Laboratoire d'Aérodologie, Université de Toulouse, CNRS, UPS, France

Brice Barret, Laboratoire d'Aérodologie, Université de Toulouse, CNRS, UPS, France

Anne Boynard, LATMOS/IPSL, UPMC Univ. Paris 06 Sorbonne Universités, UVSQ, CNRS, Paris, France; SPASCI, Ramonville Saint-Agne, 31520, France

John Burrow, Institute of Environmental Physics, University of Bremen, Germany

Catherine Clerbaux, LATMOS/IPSL, UPMC Univ. Paris 06 Sorbonne Universités, UVSQ, CNRS, Paris, France

Guanyu Huang, Harvard-Smithsonian Center for Astrophysics, Cambridge, Massachusetts, USA

Brian Kerridge, Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire, UK

Barry Latter, Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire, UK

Xiong Liu, Harvard-Smithsonian Center for Astrophysics, Cambridge, Massachusetts, USA

Nabiz Rahpoe, Institute of Environmental Physics, University of Bremen, Germany

Alexei Rozanov, Institute of Environmental Physics, University of Bremen, Germany

Catherine Wespes, Université libre de Bruxelles (ULB), Atmospheric Spectroscopy, Service de Chimie Quantique et Photophysique, Brussels, Belgium

Jerald Ziemke, Morgan State University, Baltimore, Maryland, USA

Abstract:

Tropospheric ozone is a pollutant detrimental to human health and crop and ecosystem productivity. Tropospheric ozone is also the third most important greenhouse gas (after CO₂ and methane), responsible for ~17% of global radiative forcing since 1750. However, the lack of a comprehensive global ozone monitoring network means that ozone's radiative forcing must be estimated by chemistry-climate models with a large error bars

of $\pm 50\%$ due to model uncertainties ($0.40 \pm 0.20 \text{ W m}^{-2}$, according to the fifth IPCC assessment report). Improvements to this estimate require an accurate observation-based quantification of the present-day tropospheric ozone burden (TOB), and greater confidence in chemistry-climate model estimates of TOB in pre-industrial times. TOB is the total mass (Tg) of ozone in the troposphere, calculated by summing all of the tropospheric column ozone (TCO) values at every point on Earth. Presently there is one published observation-based estimate of TOB, which comes from the OMI/MLS satellite instruments on NASA's Aura satellite. Recently, four new satellite products have been developed for measuring TCO and TOB, with two based on the OMI satellite instrument and two based on the IASI satellite instrument. The first intercomparison of these products will soon be published as a component of the Tropospheric Ozone Assessment Report (TOAR). While all five products show the same general tropospheric ozone features across the globe, they differ in absolute TCO quantities and they also differ in terms of decadal trends. The next step is to evaluate all products against the exact same set of in situ ozone observations to gauge the performance of each product in different regions of the world. We will present preliminary results from this evaluation which relies on daily IAGOS commercial aircraft profiles above Frankfurt, Germany and weekly NOAA GMD ozonesonde profiles above Hilo, Hawaii; Trinidad Head, California; and Boulder, Colorado.