

4.143 Important contributions of sea-salt aerosols to atmospheric bromine cycle in the Antarctic coasts.

Presenting Author:

Keiichiro Hara, Fukuoka Univ., harakei@fukuoka-u.ac.jp

Abstract:

Polar sunrise activates reactive bromine (BrO_x) cycle on the Antarctic coasts. BrO_x chemistry relates to depletion of O_3 and Hg in polar regions. Earlier studies have pointed out “blowing snow” as a source of atmospheric BrO_x . However, surface O_3 depletion and BrO enhancement occurs rarely under blowing snow conditions at Syowa Station, Antarctica. Therefore, trigger processes for BrO_x activation other than the heterogeneous reactions on blowing snow particles must be considered. This study shows that enhancement of sea-salt aerosols (SSA) and heterogeneous reactions on SSA are the main key processes for atmospheric BrO_x cycle activation. Blowing snow had Br^- enrichment, in contrast to strong Br^- depletion in SSA. In-situ aerosol measurements and satellite BrO measurements demonstrated clearly that a BrO plume appeared simultaneously in SSA enhancement near the surface. Results show that surface O_3 depletion at Syowa Station occurred in aerosol enhancement because of SSA dispersion during the polar sunrise. Amounts of the depleted Br^- from SSA were matched well to the tropospheric vertical column density of BrO and BrO_x concentrations found in earlier work. Our results indicate that SSA enhancement by strong winds engenders activation of atmospheric BrO_x cycles via heterogeneous reactions on SSA.