

4.250 Source apportionment of the marine aerosol over Atlantic Ocean by combining particle chemical composition and number concentration.

Early Career Scientist

Presenting Author:

Shan Huang, Institute for Environmental and Climate Research, Jinan University, Guangzhou, Guangdong, China , shanhuang_pku@163.com

Co-Authors:

Zhijun Wu, Peking University, Beijing, China

Laurent Poulain, Leibniz Institute for Tropospheric Research, Leipzig, Germany

Wolfram Birmili, Federal Environment Agency, Berlin, Germany

Hartmut Herrmann , Leibniz Institute for Tropospheric Research, Leipzig, Germany

Alfred Wiedensohler, Leibniz Institute for Tropospheric Research, Leipzig, Germany

Abstract:

The marine aerosol plays an important role in global climate regulation and marine biogenic system. However, aerosol particles in the marine boundary layer (MBL) are seldom from exclusively marine origins. In this study, chemically distinct groups are identified by k-means clustering, in an exploratory way, on chemical composition (provided by a High Resolution Time of Flight Aerosol Mass Spectrometer, HR-ToF-AMS) and particle number concentration (PNC) of MBL aerosol particles, and linked to the air mass origins. The analysis is based on a unique dataset derived from 4 research cruises over the Atlantic Ocean in 2011 and 2012, covering spatial range from 53°S to 53°N in spring and autumn. In total 12 variables were used for clustering, including PNC of aerosol particles and 11 chemical components. With this algorithm, 4 clusters were identified based on the distinct chemical composition corresponding to 2 marine types and 2 continental types. Moreover, when compared to air mass back trajectories, each cluster can be attributed to a specific air mass type and/or influenced by location. The marine 1 cluster was characterized by dominating contribution of dimethyl sulfide (DMS) oxidation products, coincidentally with air masses from the ocean. The marine 2 cluster containing significant nitrogen products, which could be attributed to biogenic amines sources, was found mainly in autumn (especially over the Southern Atlantic) within marine air masses. The continental cluster appeared only in the Northern Hemisphere, in most case associated with air masses from the Europe. The combustion cluster occupied a wide area between 15 °S and 15 °N in autumn (November), influenced by the seasonal continental outflow (e.g. biomass burning). The resulting clusters are repeatable in the same season along the similar ship tracks, which makes the method useful for modelization on predicting physicochemical properties of MBL aerosol.